

Preprint DFPD/94/TH/57

Charge Exchange Processes between Excited Helium and Fully Stripped Ions

Luca Salasnich

Dipartimento di Fisica "Galileo Galilei" dell'Università di Padova,

Via Marzolo 8, I 35131 Padova, Italy

and

Dipartimento di Fisica dell'Università di Firenze,

INFN, Sezione di Firenze,

Largo E. Fermi 2, I 50125 Firenze, Italy

Fabio Sattin

Istituto Gas Ionizzati del CNR, Associazione Euratom-ENEA-CNR


Corso Stati Uniti 4, I 35020 Padova, Italy

Abstract

We made a classical trajectory Monte Carlo (CTMC) calculation of state selective cross sections for processes between some light ions and excited helium. The results, useful for analysis of spectroscopic data of fusion devices, are in good agreement with theoretical predictions of scaling laws.

To be published in Phys. Rev. A

Impurities in high temperature plasma devices are mostly found as fully stripped ions. No spectral line is emitted by these ions, but they may be detected through characteristic lines emitted by one-electron ions found in excited states after a process of charge exchange (CX) with a neutral atom:



where N is the neutral atom, X is the ion, n and n' are quantum numbers, and $h\nu$ is the emitted photon.

In fusion devices beams of hydrogen, helium and lithium are usually injected into the plasma to heat it or to monitor such plasma parameters as ion density, temperature, velocity of rotation [1]. Extraction of useful informations from emission measurements relies on knowledge of the effective cross section for process (1).

Process (1) may take place with N in the ground or in an excited state: charge exchange involving neutrals in the ground state predominantly populate hydrogen-like states with principal quantum number $n \leq 5$ for low- Z ions, up to oxygen [1]. The decay of these states occurs mostly through $\Delta n = 1$ transitions, corresponding to emission in the X -ray or ultraviolet region [1]. The emission in the visible region is originated from states with higher n (e.g. transitions $10 \rightarrow 9$ at 6070 Å of O^{7+} or $8 \rightarrow 7$ at 5292 Å of C^{5+}); in this spectral range the detection of radiation is simpler and multi-chord systems, to spatially resolve the emission, are usually working. Cross sections for electron capture into these states from ground state neutrals are much smaller, but, on the other hand, the contribution from excited neutrals is greatly enhanced [1] so, even if the atoms excited by Coulomb collisions

with plasma particles constitute a small fraction of the entire beam, they may be the major responsible for the emission. This implies that to deduce informations from CX emission it is necessary to have data on the processes involving excited neutrals.

In this work we propose to calculate partial cross sections for process (1) (because of their much smaller cross section double electron processes are not here considered) between fully stripped ions and helium in the first excited state. We focus on conditions frequently encountered in plasma devices : the collision energy is varied in the range below 110 KeV/amu and the ionic charge varies from 3 to 8.

A fully quantum mechanical treatment of the scattering process for such a system would be difficult because of the large number of states involved. The classical trajectory Monte Carlo (CTMC) method , where the quantum system is replaced by a classical one with all the particles interacting through Coulomb forces and their motion as governed by Newton's law, has proven to be a feasible and reliable tool in this kind of calculations.

The system helium+ion actually represents a four-body problem, but the excited electron of helium plays a major role, so we chose to substitute the other electron and the nucleus altogether with a point-like particle interacting with the other charges through a central model potential [2], which in atomic units (a.u.) can be written:

$$V_{mod}(r) = -\frac{1}{r} - \frac{(1 + 0.4143r)}{r} \exp(-2.499r). \quad (2)$$

The first eigenvalues of the Schrödinger equation with this potential are -0.904 a.u. and -0.156 a.u.. It may be proved, following similar lines

as in [3], that V_{mod} provides a good approximation of electron moment distribution, which is a critical requisite for every CTMC computation [3,4].

Initialization of the system was performed according to the procedure developed by Reinhold and Falcon [3]. The initial distance between nuclei was set to $10Z$, but checks were made to larger distances, up to $20Z$, to ensure independence of the results from this parameter. For each energy a number of runs ranging from $4 \cdot 10^3$ to $27 \cdot 10^3$ was made to have a reasonable statistical error (< 10 per cent for the processes characterized by the greater frequencies). We have considered ions which are of interest in fusion devices: carbon and oxygen usually constitute the most abundant impurities, while beryllium and boron are often used to condition vacuum chamber inner walls. Only distribution on the principal quantum number n was considered: at the densities and quantum numbers considered there is a complete l -mixing before spontaneous decay take place [5].

In Fig. 1 are plotted electron capture cross sections *vs* n for different energies. The position of the maximum in the n -distribution of the cross section closely follows the rule:

$$n_{max} = \frac{Z^{3/4}}{\sqrt{-2I}}, \quad (3)$$

where $I = -0.156$ a.u. is the ionization potential of the helium. Relation (3) is already known to hold for collisions between ions and H or Li beams.

If compared with analogue curves plotted in [6] relatively to collisions between ions and He in the ground state, we notice that our distributions are much broader. The spreading in n is due the increasing of the ratio v_p/v_e , where v_p is the impact velocity of ion, and v_e is the initial orbital velocity of

the electron [5].

Scaling formulas with respect to charge of the ion, principal quantum number of the neutral, and relative energy of incidence have been derived for electron capture processes involving hydrogen. We verified all our data lie on a single curve after the rescaling

$$\sigma'(E') = \frac{\sigma(E)}{n^4 Z}, \quad E' = \frac{n^2 E}{\sqrt{Z}}, \quad (4)$$

(see Fig. 2). Scaling (4) has been suggested, for instance, in [7]. Our scaling well resembles those obtained in similar calculations with H or He in the ground state (for a comparison see the curves plotted in [6]). In our case n in eq. (4) is the effective (noninteger) quantum number.

In alternative to process (1) single ionization is also a possible result:



For ionization a similar relation to (4) has been suggested in [8] to hold, as a result of a close-coupling calculation on ground state and excited lithium:

$$\sigma'_{ion}(E') = \frac{\sigma_{ion}(E)}{n^4 Z^{1.3}}, \quad E' = \frac{n^2 E}{\sqrt{Z}}, \quad (6)$$

Even though our plots cannot discriminate little differences in the exponent of Z , we may state that our results are very similar to those of [8]; our reduced cross sections are plotted in Fig. 3.

In order to make sure that V_{mod} may represent a good enough potential we repeated some of the calculations by using a more sophisticated potential: in Fig. 4 we plotted partial cross sections for the scattering with Beryllium at 10 KeV/amu as calculated with V_{mod} and with the model potential reported in references [9,10], the agreement being quite good.

Summarizing, we have extended a well known method of calculation to processes not previously considered, which are of importance for the diagnostics of high temperature plasmas. Our results are in good accordance with extrapolations from previous calculations and empirical laws, so we may be confident on their accuracy.

* * * * *

This work has been partially supported by the Ministero dell'Università e della Ricerca Scientifica e Tecnologica (MURST). We wish to thank Dr. M.E. Puiatti and Dr. L. Carraro for their careful reading of the manuscript.

Figure Captions

Figure 1: n -distribution for one electron capture between helium and beryllium, boron, carbon and oxygen. Cross sections are in 10^{-16} cm^2 . Dotted curve correspond to an impact energy of $E = 30 \text{ KeV/amu}$, barred curve to $E = 20 \text{ KeV/amu}$ and solid curve to $E = 10 \text{ KeV/amu}$.

Figure 2: Reduced cross section $\sigma/(n^4 Z)$ for one electron capture *vs* reduced energy $En^2/Z^{1/2}$. Cross sections are in 10^{-16} cm^2 .

Figure 3: Reduced cross section $\sigma/(n^4 Z^{1.3})$ for single ionization *vs* reduced energy $En^2/Z^{1/2}$. Cross sections are in 10^{-16} cm^2 .

Figure 4: Partial cross sections σ for single electron capture from Beryllium at 10 KeV/amu obtained through use of two different model potentials. Dashed line: V_{mod} of equation (2); solid line: model potential from [9,10]. Cross sections are in 10^{-16} cm^2 .

References

- [1] R.C. Isler: *Plasma Phys. Control. Fus.* **36**, 171 (1994)
- [2] B.H. Bransden, A.M. Ermolaev and R. Schingal: *J. Phys. B: At. Mol. Phys.* **17**, 4515 (1984)
- [3] C.O. Reinhold and C.A. Falcon: *Phys. Rev. A* **33**, 3859 (1986)
- [4] A.E. Wetmore and R.E. Olson: *Phys. Rev. A* **38**, 5563 (1988)
- [5] R.J. Fonk, D.S. Arrow and P.K. Jaehnig: *Phys. Rev. A* **29**, 3388 (1984)
- [6] L. Meng, C.O. Reinhold and R.E. Olson: *Phys. Rev. A* **42**, 5286 (1990)
- [7] R.E. Olson: *J. Phys. B: At. Mol. Phys.* **13**, 483 (1980)
- [8] J. Schweinzer, D. Wutte and H.P. Winter: *J. Phys. B: At. Mol. Opt. Phys.* **27**, 137 (1994)
- [9] G. Peach, S.L. Willis and M.R.C. McDowell: *J. Phys. B: At. Mol. Phys.* **18**, 3921 (1985)
- [10] S. L. Willis, G. Peach, M.R.C. McDowell and J. Banerji: *J. Phys. B: At. Mol. Phys.* **18**, 3939 (1985)